



University of HUDDERSFIELD

University of Huddersfield Repository

Feltham, P. and Andrews, Ian

On the F-Centre in Calcium Fluoride

Original Citation

Feltham, P. and Andrews, Ian (1966) On the F-Centre in Calcium Fluoride. *Physica Status Solidi*, 15. K97-K98. ISSN 0031-8957

This version is available at <http://eprints.hud.ac.uk/id/eprint/5994/>

The University Repository is a digital collection of the research output of the University, available on Open Access. Copyright and Moral Rights for the items on this site are retained by the individual author and/or other copyright owners. Users may access full items free of charge; copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational or not-for-profit purposes without prior permission or charge, provided:

- The authors, title and full bibliographic details is credited in any copy;
- A hyperlink and/or URL is included for the original metadata page; and
- The content is not changed in any way.

For more information, including our policy and submission procedure, please contact the Repository Team at: E.mailbox@hud.ac.uk.

<http://eprints.hud.ac.uk/>

short note

reprinted from

physica
status
solidi

phys. stat. sol. 15, K97 (1966)

Department of Physics, Brunel College, London

On the F-Centre in Calcium Fluoride

By

P. FELTHAM and I. ANDREWS

In a recent note Fleming (1) referred to our calculations of the F-centre absorption wavelength in calcium fluoride (2), suggesting that mathematical expediency may have led us to an incorrect choice of the lattice potential well. In the determination of the shape of the well, shown schematically as the full line 1 in Fig. 1, we utilised a scale model of the calcium fluoride lattice to find the approximate shape of curve 1 in the important volume of high charge density, contained within the fluorine vacancy, i.e. extending from the origin up to about r_F in Fig. 1. On going in the [111] direction singularities occur on the well surface at sites of the calcium and fluorine "point" ions, a section through the well is represented by the dashed curve.

The problem of determining a well shape suitable for continuum treatments of colour-centre absorption wavelengths is well known (3, 4) and we did not consider it necessary to indicate or discuss the discontinuities, here shown on the dashed curve, specifically, depicting only the part lying between the origin and the nearest calcium ion, corresponding to the branch denoted by curve 2 in Fig. 1.




Fig. 1

Fleming appears to be under the erroneous impression that this branch of the dashed curve should be taken as the lattice potential rather than our curve 1. It is obvious that such a mistaken assumption must lead to a gross underestimate of the 1s-2p transition energy of the F-electron, as, in fact, he finds to be the case.

References

- (1) R.J. FLEMING, phys. stat. sol. 13, K87 (1966).
- (2) P. FELTHAM and I. ANDREWS, phys. stat. sol. 10, 203 (1965).
- (3) S.R. TIBBS, Trans. Faraday Soc. 35, 1471 (1936).
- (4) J.H. SIMPSON, Proc. Roy. Soc. A 197, 269 (1949).

(Received April 21, 1966)