

A Review of Evidence for Negative Pressure Wound Therapy (NPWT) use Post Spinal Surgery



Ross A. Atkinson¹
PhD, Spinal Research
Coordinator

Karen J. Ousey²
PhD, RGN, Reader,

Steve Lui² MSc,
Senior Lecturer,

John Bradley Williamson³
MChB, FRCS,
Consultant in
Spinal Surgery

¹Greater Manchester
Neurosciences Centre;
and Honorary Research
Associate – The University
of Manchester,
Manchester Academic
Health Science Centre,
Salford Royal NHS
Foundation Trust,
C235 Clinical Sciences
Building, Stott Lane,
Salford, M6 8HD.

²School of Human and
Health Sciences,
Centre for Health and
Social Care Research,
University of Huddersfield,
Queensgate, Huddersfield,
West Yorkshire, HD1 3DH.

³Salford Royal NHS
Foundation Trust,
Stott Lane,
Salford, M6 8HD

Correspondence:
ross.atkinson@nhs.net

Conflict of interest: none

EXTENDED ABSTRACT

Aims

To systematically search, critically appraise and summarise randomised controlled trials (RCTs) and non-RCTs assessing the effectiveness of negative pressure wound therapy (NPWT) in patients with a surgical spinal wound.

Methods

A systematic review based on search strategies recommended by the Cochrane Back and Wounds Review Groups was undertaken using the Cochrane Library, MEDLINE, EMBASE and CINAHL databases. Any publications between 1950 and 2011 were included. Funding to undertake this systematic review was received from the University of Huddersfield Collaborative Venture Fund and KCI Medical.

Results

Nine retrospective studies⁽¹⁻⁹⁾ and five case studies⁽¹⁰⁻¹⁴⁾ of patients with spinal wound complications were included in this systematic review. No RCTs were found. Only one study described more than 50 patients⁽⁴⁾. Generally, a pressure of -100 to -125 mmHg was used in adult patients^(1,8,12).

Duration of NPWT in situ ranged from three to 186 days^(2,5,6,8,13). Wound healing was assessed every two to three days and generally completed between seven days and 16 months^(1-5,9-14). NPWT is contraindicated in the presence of active cerebrospinal fluid leak⁽¹⁾, metastatic or neoplastic disease in the wound^(9,10), in patients with an allergy to the NPWT dressing⁽⁹⁾, and in those with a bleeding diathesis⁽¹⁾.

Discussion

We identified no RCTs discussing the use of NPWT in the management of surgical spinal wounds, and limited low quality evidence demonstrating that NPWT can be used effectively in this type of patient. In an RCT in obese patients undergoing total knee arthroplasty, no difference in the time taken to achieve a dry wound with NPWT as compared with a sterile gauze has been reported⁽¹⁵⁾. Importantly, that study was terminated early due to the presence of skin blisters associated with the NPWT dressing; an adverse effect which has not been reported in the spine literature. Furthermore, Dorafshar and colleagues⁽¹⁶⁾ concluded that NPWT did not provide superior

References

- Jones GA, Butler J, Lieberman I, Schlenk R. Negative-pressure wound therapy in the treatment of complex postoperative spinal wound infections: complications and lessons learned using vacuum-assisted closure. *J Neurosurg Spine* 2007; 6(5):407-411.
- Labler L, Keel M, Trentz O, Heinzlmann M. Wound conditioning by vacuum assisted closure (V.A.C.) in postoperative infections after dorsal spine surgery. *Eur Spine J* 2006; 15(9):388-396.
- Mehbod AA, Ogilvie JW, Pinto MR, Schwender JD, Transfeldt EE, Wood KB et al. Postoperative deep wound infections in adults after spinal fusion: management with vacuum-assisted wound closure. *J Spinal Disord Tech* 2005; 18(1):14-17.
- Ploumis A, Mehbod AA, Dressel TD, Dykes DC, Transfeldt EE, Lonstein JE. Therapy of spinal wound infections using vacuum-assisted wound closure: risk factors leading to resistance to treatment. *J Spinal Disord Tech* 2008; 21(5):320-323.
- Antony S, Terrazas S. A retrospective study: clinical experience using vacuum-assisted closure in the treatment of wounds. *J Natl Med Assoc* 2004; 96(8):1073-1077.
- Canavese F, Gupta S, Krajchich JI, Emara KM. Vacuum-assisted closure for deep infection after spinal instrumentation for scoliosis. *J Bone Joint Surg Br* 2008; 90(3):377-381.
- Canavese F, Gupta S, Emara KM, Krajchich JI. Use of the vacuum assisted closure in instrumented spinal deformities for children with neuromuscular scoliosis who developed post-operative deep spinal infection. *Developmental Medicine and Child Neurology* 2009; 51(5):50.
- Horn PL, Ruth B, Kean JR. Use of wound V.A.C. therapy in pediatric patients with infected spinal wounds: a retrospective review. *Orthop Nurs* 2007; 26(5):317-322; quiz 323-324.
- Zehnder SW, Place HM. Vacuum-assisted wound closure in postoperative spinal wound infection. *Orthopedics* 2007; 30(4):267-272.
- Yuan-Innes MJ, Temple CL, Lacey MS. Vacuum-assisted wound closure: a new approach to spinal wounds with exposed hardware. *Spine (Phila Pa 1976)* 2001; 26(3):E30-33.
- Adams J, Hakim EM. A critical analysis of existing evidence on the management of a wound with Malignant Desmoplastic Melanoma and Neurofibromatosis: a case report. *Rehabilitation Oncology* 2009; 27(1):26-27.
- van Rhee MA, de Klerk LW, Verhaar JA. Vacuum-assisted wound closure of deep infections after instrumented spinal fusion in six children with neuromuscular scoliosis. *Spine J* 2007; 7(5):596-600.
- Vicario C, de Juan J, Esclarin A, Alcobendas M. Treatment of deep wound infections after spinal fusion with a vacuum-assisted device in patients with spinal cord injury. *Acta Orthop Belg* 2007; 73(1):102-106.



EWMA 2012

outcomes with regards to wound volume and surface area over wall suction applied to a sealed gauze dressing in patients with acute wounds resulting from trauma, dehiscence or surgery. However, a further study in knee arthroplasty patients found that NPWT was associated with a reduction in post-operative development of seromas and improved healing⁽¹⁷⁾.

Conclusions

The evidence for NPWT use in orthopaedic surgery is ambiguous, and at best low quality in spine patients. Published reports are limited to small retrospective and case studies, with no reports of NPWT being used as a prophylactic treatment in spinal patients. Larger, prospective RCTs of NPWT are needed to support the current evidence that it is effective in treating spinal wound complications. In addition, future studies should investigate the use of NPWT as a prophylactic treatment to promote wound healing and prevent infection, and report data relating to safety and health economics. ■

14. Hwang JH, Modi HN, Suh SW, Hong JY, Yang JH, Park JH. Maggot debridement therapy for postsurgical wound infection in scoliosis: a case series in five patients. *Spine (Phila Pa 1976)* 2011; 36(4):313-319.
15. Howell RD, Hadley S, Strauss E, Pelham FR. Blister formation with negative pressure dressings after total knee arthroplasty. *Current Orthopaedic Practice* 2011; 22(2):176-179.
16. Dorafshar AH, Franczyk M, Gottlieb LJ, Wroblewski KE, Lohman RF. A prospective randomized trial comparing subatmospheric wound therapy with a sealed gauze dressing and the standard vacuum-assisted closure device. *Annals of Plastic Surgery* 2011; 69(1):79-84.
17. Pachowsky M, Gusinde J, Klein A, Lehl S, Schulz-Drost S, Schlechtweg P, et al. Negative pressure wound therapy to prevent seromas and treat surgical incisions after total hip arthroplasty. *International Orthopaedics* 2011; 36(4):719-722.

THE EWMA UNIVERSITY CONFERENCE MODEL (UCM) in Copenhagen



The EWMA UCM programme offers students of wound management from institutes of higher education across Europe the opportunity to take part of their academic studies whilst participating in the EWMA Conference.

The opportunity of participating in the EWMA UCM is available to all teaching institutions with wound management courses for health professionals.

The UCM programme at the EWMA 2013 Conference in Copenhagen will offer networking opportunities between the students from various UCM groups, UCM Lectures as well as assignments and workshops arranged specifically for the UCM students.

EWMA strongly encourages teaching institutions and students from all countries to benefit from the possibilities of international networking and access to lectures by many of the most experienced wound management experts in the world.

Yours sincerely

Zena Moore,
Chair of the Education Committee, Immediate Past President

Participating institutions:



Donau Universität Krems
Austria



Haute École de Santé
Geneva, Switzerland



HUB Brussels
Belgium



Lithuanian University of Health Sciences
Lithuania

PROFESSIONSHØJSKOLEN



Metropolitan University College
Denmark



University of Hertfordshire
United Kingdom



Universidade Católica Portuguesa
Porto, Portugal

For further information about the EWMA UCM, please visit the Education section of the EWMA website www.ewma.org or contact the EWMA Secretariat at ewma@ewma.org