



University of HUDDERSFIELD

University of Huddersfield Repository

Vanin, Stefano, Bonizzoli, Manuela, Migliaccio, Maria Luisa, Buoninsegni, Laura Tadini, Bugelli, Valentina, Pinchi, Vilma and Focardi, Martina

A case of insect colonization before the death

Original Citation

Vanin, Stefano, Bonizzoli, Manuela, Migliaccio, Maria Luisa, Buoninsegni, Laura Tadini, Bugelli, Valentina, Pinchi, Vilma and Focardi, Martina (2017) A case of insect colonization before the death. *Journal of Forensic Sciences*. ISSN 0022-1198

This version is available at <http://eprints.hud.ac.uk/id/eprint/31955/>

The University Repository is a digital collection of the research output of the University, available on Open Access. Copyright and Moral Rights for the items on this site are retained by the individual author and/or other copyright owners. Users may access full items free of charge; copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational or not-for-profit purposes without prior permission or charge, provided:

- The authors, title and full bibliographic details is credited in any copy;
- A hyperlink and/or URL is included for the original metadata page; and
- The content is not changed in any way.

For more information, including our policy and submission procedure, please contact the Repository Team at: E.mailbox@hud.ac.uk.

<http://eprints.hud.ac.uk/>

A case of insect colonization before the death

Stefano Vanin, PhD¹, Manuela Bonizzoli, MD³, Marialuisa Migliaccio, MD³, Laura Tadini

Buoninsegni, MD³, Valentina Bugelli, MD², Vilma Pinchi, PhD, DDS², Martina Focardi, MD²

¹School of Applied Sciences, University of Huddersfield, UK; ²Department of Health Sciences, Forensic Medicine Section, University of Florence, Italy; ³Department of Neuromusculoskeletal and Sensory Organs, University of Florence, Italy

ABSTRACT

Forensic entomology is a branch of forensic science in which insects are used as evidence in legal investigations relating to humans, domestic animals and wildlife. One of the theoretical pillars on which the discipline is based concerns the fact that flies colonize a body after death. However in cases of myiasis maggots are present before death, with consequences in the correct estimation of the minimum Post Mortem Interval (mPMI).

We report here the case of a woman, largely colonized by fly larvae, who has lain alive in her garden for four days prior to being rescued. Larvae were found on the conjunctivae, the bronchi, the rectum and vagina. The woman's death, two months later, was caused by tetanus.

The consequences of myiasis on mPMI estimation are here discussed. In fact, despite she was still alive larvae, showed an estimated age of 1.5-2.5 days, based on environmental and body temperature.

Key-Words: forensic science, forensic entomology, forensic pathology, PMI estimation, myiasis, Calliphoridae

Forensic entomology is the branch of forensic sciences in which insects are used as evidence in legal investigations related to humans, domestic animals and wildlife (1-3). The examination, identification and analysis of the insects associated with human remains, combined with the

knowledge of insect biology and the meteorological parameters, especially temperature, can provide a further level of detail in addition to medical and anthropological data in reconstructing the events that occurred close to the death.

In particular, necrophagous insects are useful for the minimum Post-Mortem Interval (mPMI) estimation, the movement of the body after the death, for the detection of drugs or poisons if soft tissues are no longer available or too decomposed (1,4). Furthermore, insects found on a crime scene can be used for human DNA extraction for identification purposes, when no suitable tissues are available (5) or even no body was found on the crime scene because removed before the police arrival (eg.: post mortem body movement).

One of the theoretical pillars on which the discipline is based, concerns the fact that flies colonize a body after the death and that the age of the insects can be estimated considering the temperature at which these organisms were exposed during their development. The estimation of the age of the first body colonizers allows the estimation of the so called “time of colonization”, commonly considered as the mPMI (6).

However in cases of myiasis, the infestation of fly larvae on living or necrotic tissues in vertebrates, maggots are already present on the body of the victim before his/her death.

Human myiasis can be classified into two groups: specific myiasis produced by flies that need a live host to develop their immature stages, and semispecific myiasis caused by flies that typically develop in decomposing organic matter but occasionally develop on/in living organisms, mostly in the presence of wounds.

The most common semispecific myiasis-causing species belong to the genera *Calliphora*, *Lucilia* (Diptera, Calliphoridae) and *Sarcophaga* (Diptera, Sarcophagidae). In humans, these species mainly develop in necrotic tissue, in cutaneous wounds with non-necrotic tissue, and in body cavities with an accumulation of secretions and excreta with a high bacterial load that constitutes the source of food for the larvae (7- 9).

In a case of myiasis, female flies can lay eggs on living subject in cases of poor personal hygiene as recorded in several homeless persons, in presence of open wounds and gangrene like in diabetes patients (9), and in case of non-conscious or non-self-sufficient/dependent persons such as cases of abandoned children and neglected elders (10).

However, in cases of poor personal hygiene or abandoned children and neglected elderly, the estimation of the colonization time can provide useful information about when the condition, allowing flies to lay eggs took place. This information can have a forensic interest in case of people neglected or in nosocomial myiasis in order to define legal responsibilities (9). In other cases like in the case here described, the colonization time can provide information about when the accident, responsible of the loss of consciousness of the “victim”, took place.

However, in all the cases previously described, the time of colonization do not correspond to the mPMI, if the subject was found already dead. These circumstances may result in relevant and critical errors on the estimation of mPMI with investigative consequences.

In this paper is presented the case of a woman, in an unconscious state and largely colonized by fly larvae, who had lain alive in her garden for several days before being rescued.

Case

In July 2015 an 84 year old woman affected by hypertension and hyperthyroidism, was found unconscious supine in the garden of her house in the suburbs of Florence (Central Italy) by the Emergency Rescue Team. The woman was wearing an undershirt and underwear. Neighbours reported that she had been seen 5 days before her rescue. Medical observations carried out during the pre-hospital evaluation indicated that the patient was hypertensive, hypoglycemic and with a strong larvae infestation on the face, especially on the conjunctivae, the nasal choanae, the mouth and the external auricular canals. An ulceration was detected in the sacral region, potentially related with her lying position. This injury was later treated with escaectomy and debridement. External examination revealed the absence of traumatic injuries, however underlined the presence of

sunburns I and II degree on exposed parts of the abdomen and thorax. An initial total body CT scan showed a left cerebral intraparenchymal-capsulo-lenticular hemorrhage causing only the compression of the left lateral ventricle. No other traumatic injuries were detected. The observed hemorrhage can be identified as the cause of the woman fall and loss of consciousness. Further investigation, based on fibrescopy, revealed that larvae were present in the bronchi, rectum, vagina and in the external auricular canals. In addition, it was observed that larval feeding activity was also responsible of multiple bilateral perforations of the tympanic membranes. A second CT scan reported a bilateral purulent ethmoidal-maxillary sinusitis that underwent a surgical drainage. No larvae were found in the affected sinuses. During the hospitalization a bacterial pneumonia was detected. During the neurological rehabilitation muscular spasms appeared and the woman died two months later. The cause of death was determined as tetanus based on the presence of trismus, stiffness in the joints and the column followed by immunologic tests. It is worth mentioning that at the Hospital admission the woman was treated with an emergency booster injection of tetanus toxoid.

Larval samples were collected from different body regions, killed in hot water, fixed in 70% Ethanol and sent to an entomologist following the standard and guidelines of the European Association for Forensic Entomology (1). Larvae were diaphanized using a hot solution of NaOH and all the entomological observations were performed with a Leica M60 stereomicroscope and a Leica DME compound. Larvae, on the II/III instars were identified as members of the Calliphoridae family. The oldest specimens showed posterior spiracles and cephaloskeleton sclerites of both the instars present at the same time, indicating the molting stage. Diagnostic features related to the shape of the oral sclerites allow the identification of the species as *Lucilia* cfr *sericata* (Meigen, 1826). (Diptera, Calliphoridae) Temperature obtained from the closest meteorological station collected during the days before the body rescue indicated a temperature average of 28 °C with a minimum of 22 and a maximum of 35 °C. This late temperature is close to the body average temperature and for this reason is more accurate in the estimation of the development in case of

internal myiasis. The larval age was estimated being between 1.5 and 2.5 days using the data published by Grassberger & Reiter (11).

Lucilia sericata is a cosmopolitan species and one of the most common body colonizers especially in the warmer seasons. This species has been reported from several outdoors Italian cases (12-14) and in a general survey carried out in 2010 on over 200 cases from Northern and Central Italy the species was found in more than 30% of the cases (15).

Lucilia sericata, as other species of the genus *Lucilia* (eg. *Lucilia cuprina* (Wiedemann, 1830), *Lucilia silvarum* Meigen, 1826, *Lucilia bufonifera* Moniez, 1876 and *Lucilia illustris* Meigen, 1826, is known as a myiasigenic species both in human and other vertebrates (16-20).

Developmental data have been produced by several authors in a range between 10 and 34 °C (for a general summery see Amendt et al (1)).

Conclusion

It is worth mentioning that, in cases like this, a later arrival of the rescue team would result with the finding of a dead woman. The estimation of the time since death based on the entomological and on the tanatho-chronological approaches clearly would give different results. For the first one the estimation would be of 1.5-2.5 days whereas applying the methods commonly used by the forensic pathologists (eg.: *algor*, *livor*, *rigor mortis*, electrolyte concentration in *vitreus humor*) the estimation would be of some hours. In fact this case clearly demonstrates that in case of dead body the time of insect colonization could not correspond to the minimum time since death. For this reason a close collaboration between forensic pathologist and forensic entomologist and a careful analysis of all possible variables are necessary, in order to answer at the investigative question about the time since death. In particular, in suspicious conditions, especially when the body is fresh, histological analysis of the colonized tissues can be helpful in order to evaluate if the colonization started before or after the death of the victim (21) and to evaluate the time of a possible agony. A

multidisciplinary approach has to be considered fundamental for a proper reconstruction of the perimortem events and for the mPMI estimation.

Acknowledgments

Authors want to thank Prof. GA Norelli, Prof. A. Peris and Dr M. Mangini for the information provided and the critical review of the manuscript. The authors are grateful to the anonymous reviewers and the editor for their suggestions.

References

1. Amendt J, Campobasso CP, Gaudry E, Reiter C, LeBlanc H, Hall MJR. Best practice in forensic entomology-standards and guidelines *Int J Legal Med.* 2007 Mar;121 (2):90–104.
2. Anderson GS. Wildlife forensic entomology: determining time of death in two illegally killed black bear cubs. *J Forensic Sci.* 1999 Jul;44(4):856-9.
3. Sanford MR, Forensic entomology of decomposing humans and their decomposing pets. *Forensic Sci Int.* 2015 Feb;247:e11-7.
4. Smith KGV. A manual of forensic entomology. Trustees of the British Museum (Natural History). Cornell University Press 1986.
5. Marchetti D, Arena E, Boschi I, Vanin S. Human DNA extraction from empty puparia. *Forensic Sci Int.* 2013 Jun 10;229(1-3):e26-9.
6. Villet MH, Cameron SR, John M. Contemporary precision, bias and accuracy of minimum post-mortem intervals estimated using development of carrion-feeding insects. In: Amendt J, Goff ML, Campobasso CP, Grassberger M. (Eds.) *Current Concepts in Forensic Entomology.* Springer 2010 Edition, 109-137.
7. Leclercq M. Les myiases. *Annales Soc Ent Fr* 1990 July;26(3):335-350.

8. Goff M.L, Campobasso C.P, Gherardi M. Forensic Implications of Myiasis. In Amendt J, Goff ML, Campobasso CP, Grassberger M. (Eds.) Current Concepts in Forensic Entomology. Springer 2010 Edition, 313-325.
9. Dutto M, Pellegrino M, Vanin S. Nosocomial myiasis in a patient with diabetes. J Hosp Infect. 2013 Jan;83(1):74-6.
10. Benecke M, Lessig R. Child neglect and forensic entomology. Forensic Sci Int. 2001 Aug 15;120(1-2):155-9.
11. Grassberger M, Reiter C. Effect of temperature on *Lucilia sericata* (Diptera: Calliphoridae) development with special reference to the isomegalen- and isomorphen-diagram. Forensic Sci Int. 2001 Aug 15;120(1-2):32-6.
12. Bugelli V, Forni D, Bassi L.A, Di Paolo M, Marra D, Lenzi S, et al. Forensic entomology and the estimation of the minimum time since death in indoor cases. J Forensic Sci. 2015 Mar;60(2):525-31.
13. Vanin S, Tasinato P, Ducolin G, Terranova C, Zancaner S, Montisci M, et al. Use of *Lucilia* species for forensic investigations in Southern Europe Forensic Sci Int. 2008 May 2;177(1):37-41.
14. Turchetto M, Lafisca S, Costantini G. Postmortem interval (PMI) determined by study sarcophagous biocenoses: three cases from the province of Venice (Italy). Forensic Sci Int. 2001 Aug 15;120(1-2):28-31.
15. Vanin S., Gherardi M. 2010. A review of 200 forensic cases involving insects from central and Northern Italy. In: Proceeding of the 7th annual meeting of the European Association for forensic Entomology (EAFE) Murcia, Spain.
16. Pezzi M, Whitmore D, Chicca M, Lanfredi M, Leis M. Traumatic myiasis caused by an association of *Sarcophaga tibialis* (Diptera: Sarcophagidae) and *Lucilia sericata* (Diptera: Calliphoridae) in a domestic cat in Italy. Korean J Parasitol. 2015 Aug;53(4):471-5.

17. Azevedo WT, Figueiredo AL, Carvalho RP, Lemos GA, Silva PF, Miranda TA, et al. Record of the First Cases of Human Myiasis by *Lucilia cuprina* (Diptera: Calliphoridae), Rio de Janeiro, Brazil. J Med Entomol 2015 Nov; 52(6):1368-73.
18. Demirel-Kaya F, Orkun Ö, Çakmak A, İnkaya AÇ, Öcal M, Erguven S. A Case of extensive wound myiasis caused by *Lucilia sericata* (Diptera: Calliphoridae) in a patient with maxillary sinus squamous cell carcinoma, in Turkey. J Arthropod Borne Dis. 2016 Jan 5;10(2):267-70.
19. Ahadzadeh EN, Ketchum HR, Wheeler R. Human Cutaneous Myiasis by the Australian Sheep Blowfly, *Lucilia cuprina* (Diptera: Calliphoridae), in Oklahoma. J Forensic Sci. 2015 Jul;60(4):1099-100.
20. Choi W, Kim GE, Park SH, Shin SE, Park JH, Yoon KC. First report of external ophthalmomyiasis caused by *Lucilia sericata* Meigen in a healthy patient without predisposing risk factors. Parasitol Int. 2015 Oct;64(5):281-3.
21. Bonelli A, Bacci S, Vannelli GB, Norelli GA. Immunohistochemical localization of mast cells as a tool for the discrimination of vital and postmortem lesions. Int J Legal Med. 2003 Feb;117(1):14-8.