



University of HUDDERSFIELD

University of Huddersfield Repository

Palit, V and Gavin, Helen

Prognostic indicators & overall survival in men presenting with PSA levels greater than 1000ng/ml.

Original Citation

Palit, V and Gavin, Helen (2010) Prognostic indicators & overall survival in men presenting with PSA levels greater than 1000ng/ml. In: British Association of Urological Surgeons Annual Meeting, 21-24 June 2010, Manchester Central. (Submitted)

This version is available at <http://eprints.hud.ac.uk/id/eprint/6780/>

The University Repository is a digital collection of the research output of the University, available on Open Access. Copyright and Moral Rights for the items on this site are retained by the individual author and/or other copyright owners. Users may access full items free of charge; copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational or not-for-profit purposes without prior permission or charge, provided:

- The authors, title and full bibliographic details is credited in any copy;
- A hyperlink and/or URL is included for the original metadata page; and
- The content is not changed in any way.

For more information, including our policy and submission procedure, please contact the Repository Team at: E.mailbox@hud.ac.uk.

<http://eprints.hud.ac.uk/>

Prognostic indicators & overall survival in men presenting with PSA levels greater than 1000ng/ml.

Authors- Victor Palit, George Delves, Mike Kimuli , Karol Rowawski , Mike Stower & Helen Gavin

Presenter- Victor Palit

Introduction- Men presenting with serum PSA above 400ng/ml have high incidence of disease related morbidity with a poor overall survival. However some men with very high PSA (>1000ng/ml) have good hormone response with survival over 5 years.

The aim of this study was to determine the PSA response to hormonal therapy and assess serum levels of alkaline phosphatase (SAP), & bone metastasis with overall survival.

Material & Methods- In this retrospective study, 91 men with prostate cancer whose PSA>1000ng/ml at initial presentation were identified between May 2004-August 2009. Data regarding the treatment, clinical and biochemical response, survival were obtained from case notes.

Results-

An independent samples t-test to examine the effect of anti-androgenic therapy on survival showed a significant difference ($t = 5.874$ $p < 0.01$, unequal variances) , suggesting a positive impact of therapy. Patients without metastasis achieved a larger fall in PSA than those with metastasis, indicating that change in PSA levels may be a good prognostic indicator. ($F_{2,12} = 4.79$, $p < 0.05$).

Chi-square test showed a significant association between SAP and metastasis status ($\chi^2 = 6.526$, $p < 0.05$) suggesting SAP levels could be used as a predictor of metastasis. Kaplan-Meier analysis showed a significant survival advantage with anti-androgenic therapy. When metastasis is factored in as a stratifying function, the distinctions become clearer. Adding the SAP levels, demonstrates a further impact.

Conclusion-Patients presenting with PSA> 1000ng/ml have poor survival. Change in PSA levels, SAP & bone metastasis could be used as a prognostic indicator for metastasis. SAP & bone metastasis could predict survival.