



# *University of* **HUDDERSFIELD**

## **University of Huddersfield Repository**

Kola, Susanna, Walsh, Jane C., Hughes, Brian M. and Howard, Siobhan

Attention focus, trait anxiety and pain perception in patients undergoing colposcopy

### **Original Citation**

Kola, Susanna, Walsh, Jane C., Hughes, Brian M. and Howard, Siobhan (2010) Attention focus, trait anxiety and pain perception in patients undergoing colposcopy. In: STAR 2010: 31st World Conference on Stress and Anxiety Research, 4th - 6th August 2010, Galway, Ireland. (Unpublished)

This version is available at <https://eprints.hud.ac.uk/id/eprint/9740/>

The University Repository is a digital collection of the research output of the University, available on Open Access. Copyright and Moral Rights for the items on this site are retained by the individual author and/or other copyright owners. Users may access full items free of charge; copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational or not-for-profit purposes without prior permission or charge, provided:

- The authors, title and full bibliographic details is credited in any copy;
- A hyperlink and/or URL is included for the original metadata page; and
- The content is not changed in any way.

For more information, including our policy and submission procedure, please contact the Repository Team at: [E.mailbox@hud.ac.uk](mailto:E.mailbox@hud.ac.uk).

<http://eprints.hud.ac.uk/>

# Attention focus, trait anxiety and pain perception in patients undergoing colposcopy

**Susanna Kola, Jane C. Walsh, Brian M. Hughes & Siobhán Howard**

31<sup>st</sup> Annual STAR Conference

August 2010



**NUI Galway**  
**OÉ Gaillimh**

School of Psychology

# Introduction

- Invasive medical procedures are considered very stressful and anxiety evoking by patients.
- Anxiety in turn affects acute pain, e.g., heightened anxiety reduces pain threshold/tolerance, increases pain intensity
- Attention is also linked to pain perception
  - Distraction
  - Sensory focusing



# Introduction

- Focus of attention a mediational role in the perception of pain - can lead to increased or decreased pain reports
- **Distraction**
  - Reduced pain perception (e.g., Devine & Spanos, 1990)
  - Increased pain tolerance (e.g. Piira et al., 2005; James & Hardardottir, 2002)
- **Sensory Focus**
  - Reduced pain perception (e.g., Ahles et al., 1983)
  - Increased pain tolerance (e.g., Keogh & Herdenfeldt, 2002)



# Introduction

- Effects of anxiety due to underlying anxious trait?
  - Failure of previous studies to account for anxious disposition
- Expectations of pain correlated with pain experience
  - Particularly important in medical situations?



# Methodology

- Aim: Evaluating two attention-based interventions, while assessing the effect of trait anxiety and pain expectancy on outcomes (self-reported pain, anxiety, and affect)
  - Colposcopy: diagnostic procedure aimed at detecting pre-cancerous cervical lesions
- 123 patients underwent procedure in one of 3 conditions:
  - Active distraction: engaged in cognitive task on laptop
  - Sensory Focus: focus on sensations in non-emotional way
  - Undirected control: standard care



# Results

- Pain expectancy was significantly related to experienced pain intensity, sensory pain, affective pain, and peak pain
  - greater expectations of pain prior to colposcopy resulted in greater self-reported pain
- Trait anxiety was not significantly related to any of the pain measures, except the McGill Pain Questionnaire item ‘miscellaneous pain’
  - Higher trait anxiety was associated with greater miscellaneous pain



# Results

- Elevated pre-colposcopy state anxiety and negative affect was associated with over-prediction of pain intensity
- Under-prediction of pain intensity was associated with greater post-colposcopy state anxiety
- However, if control for experienced pain intensity, this association is no longer significant
  - Thus, heightened state anxiety following colposcopy not due to under-prediction of pain, but rather experienced pain





# Discussion

- No observed differences in self-reported pain perception between the three groups, while controlling for pain expectancy and trait anxiety
  - Active distraction and sensory focus patients reported similar pain ratings to women in the control condition
- Pain expectancy found to be associated with experienced pain
  - Pain expectancy should thus be considered in future studies
- Trait anxiety did not have an effect on pain perception



# Conclusion

- Interventions aimed at producing accurate pain expectations may lead to reductions in pain
- Trait anxiety does not appear to directly influence pain perception, but indirectly through elevations in state anxiety and negative affect
  - Anxious state thus a better predictor for identifying colposcopy patients at risk for experiencing pain during the examination



# Thank you!

Susanna Kola is now at the Department of Behavioural  
and Social Sciences at the University of Huddersfield

For further information email [s.kola@hud.ac.uk](mailto:s.kola@hud.ac.uk)



NUI Galway  
OÉ Gaillimh

School of Psychology