



University of HUDDERSFIELD

University of Huddersfield Repository

Kola, Susanna, Walsh, Jane C., O'Gorman, Tom and Lee, John

The role of attention in reducing anxiety, pain and distress associated with a stressful diagnostic procedure

Original Citation

Kola, Susanna, Walsh, Jane C., O'Gorman, Tom and Lee, John (2006) The role of attention in reducing anxiety, pain and distress associated with a stressful diagnostic procedure. In: 27th International Conference of the Stress and Anxiety Research Society, July 2006, Crete. (Unpublished)

This version is available at <http://eprints.hud.ac.uk/id/eprint/9764/>

The University Repository is a digital collection of the research output of the University, available on Open Access. Copyright and Moral Rights for the items on this site are retained by the individual author and/or other copyright owners. Users may access full items free of charge; copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational or not-for-profit purposes without prior permission or charge, provided:

- The authors, title and full bibliographic details is credited in any copy;
- A hyperlink and/or URL is included for the original metadata page; and
- The content is not changed in any way.

For more information, including our policy and submission procedure, please contact the Repository Team at: E.mailbox@hud.ac.uk.

<http://eprints.hud.ac.uk/>

The role of attention in reducing anxiety, pain and distress associated with a stressful diagnostic procedure

Susanna Kola, Jane Walsh,
Tom O’Gorman & John Lee





Invasive Medical Procedures

- Are perceived by most patients as highly stressful and anxiety provoking
- Heightened anxiety levels can lead to a variety of complications, incl. heightened experience of pain
- Flexible sigmoidoscopy is an invasive diagnostic procedure to detect bowel abnormalities, and is routinely performed without sedation or analgesia despite being uncomfortable and painful



Pain

- An unpleasant sensory or emotional experience, associated with actual or potential tissue damage or described in terms of such damage (IASP, 1979)
- Pain experience results from integration of sensory-discriminative, affective-emotional, and cognitive-evaluative axes (Melzack & Katz, 1994)



Anxiety and Pain

- Anxiety is a future-oriented emotion, characterised by negative affect and apprehensive anticipation of potential threat

- State anxiety is associated with
 - Reduced pain tolerance (Carter et al., 2002)
 - Increased pain perception (Williams, 1999; Jones et al., 2002)
 - Prolongation of pain experience (Williams, 1999)
 - Lowered pain threshold (Williams, 1999; Michelotti et al., 2000)

- Trait anxiety less studied, but
 - HTA ↓ pain tolerance (James & Hardardottir, 2002)
 - HTA males ↓ tolerance, ↑ pain intensity (Jones et al., 2003)
 - HTA ↑ pain intensity (Tang & Gibson, 2005)




Attention and Pain

- Focus of attention a mediational role in the perception of pain - can lead to increased or decreased pain reports
- **Distraction**
 - Reduced pain perception (e.g., Devine & Spanos, 1990)
 - Increased pain tolerance (e.g. Piira et al., 2005; James & Hardardottir, 2002)
- **Sensory Focus**
 - Reduced pain perception (e.g., Ahles et al., 1983)
 - Increased pain tolerance (e.g., Keogh & Herdenfeldt, 2002)



Attention and Pain

- **Limited Capacity Resource Theory of Attention** (Kahneman, 1973)
 - Attention is of limited capacity and distraction reduces the available resources to process pain stimulus
- **Parallel Processing Theory of Pain Distress** (Leventhal & Everhart, 1979)
 - Involve focusing attention on the physical sensations of pain and discomfort in a concrete, objective, non-distressing way



Distraction vs. Sensory Focus: Explanation for equivocal results

- Pain intensity or threat level (Eccleston & Crombez, 1999)
- Temporal factors (Suls & Fletcher, 1985)
- Anxiety may mediate the relationship between attention and pain (James & Hardardottir, 2002)



Study Aims

- Intervention study to reduce patient anxiety, pain and distress relating to flexible sigmoidoscopy using different cognitive attention strategies
- Investigation of which strategy is the most effective, and for whom
- Examine various psychological variables in relation to the experience of the procedure



Methodology

- Randomised controlled trial
- First time flexible sigmoidoscopy patients
- Random assignment to one of four conditions
 - Audio-Visual Cognitive Distraction – lexical decision task
 - Sensory Focus – pay close attention to sensations
 - Audio-Visual Relaxation – nature views and relaxing music
 - Control – standard care

Audio-Visual Distraction Conditions

- In the Cognitive Distraction and Relaxation conditions patients to wear virtual i-glasses
- **Cognitive Distraction** – lexical decision task
- **Relaxation** – view “DVD 'At Water's Edge..' (SereneVision Productions Inc.)



Sensory Focus and Control Conditions

- In the **Sensory Focus** condition patients are asked to pay close attention to their sensations while watching the real-time FS on the monitor
- **Control Condition** – Standard Care





Measures

<u>Pre-procedure</u>	<u>During</u>	<u>Post-procedure</u>
Demographic info	BP/HR	BP/HR
BP/HR	Behavioural	Discomfort ratings
STAI (state & trait)	observations of	Pain intensity
Marlowe-Crowne	distress	ratings
PANAS	Depth of Insertion	MPQ
FPQ-III	Length of time of	STAI (state)
Discomfort	procedure	Coping with Pain
Expectancy		Questionnaire
Pain Expectancy		PANAS
		MBSS



Predictions

- **Distraction** – greater adjustment and lower pain intensity levels
- **Sensory focus** – greater adjustment and better pain outcomes if patients find it hard to disengage attention
- Likely that individual difference variables will interact with focus of attention, especially trait anxiety (James & Hardardottir, 2002)