



# *University of* **HUDDERSFIELD**

## **University of Huddersfield Repository**

Atkinson, Ross A., Davies, Benjamin, Jones, Anna, Van Popta, Dmitri, Ousey, Karen and Stephenson, John

Survival of patients undergoing surgery for metastatic spinal tumours and the impact of surgical site 2 infection

### **Original Citation**

Atkinson, Ross A., Davies, Benjamin, Jones, Anna, Van Popta, Dmitri, Ousey, Karen and Stephenson, John (2016) Survival of patients undergoing surgery for metastatic spinal tumours and the impact of surgical site 2 infection. *Journal of Hospital Infection*, 94 (1). pp. 80-85. ISSN 0195-6701

This version is available at <https://eprints.hud.ac.uk/id/eprint/28619/>

The University Repository is a digital collection of the research output of the University, available on Open Access. Copyright and Moral Rights for the items on this site are retained by the individual author and/or other copyright owners. Users may access full items free of charge; copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational or not-for-profit purposes without prior permission or charge, provided:

- The authors, title and full bibliographic details is credited in any copy;
- A hyperlink and/or URL is included for the original metadata page; and
- The content is not changed in any way.

For more information, including our policy and submission procedure, please contact the Repository Team at: [E.mailbox@hud.ac.uk](mailto:E.mailbox@hud.ac.uk).

<http://eprints.hud.ac.uk/>

**FULL TITLE**

Survival of patients undergoing surgery for metastatic spinal tumours and the impact of surgical site infection

**RUNNING TITLE**

SSI and survival in spinal tumour patients

**AUTHORS**

**Ross A. Atkinson**<sup>1,2</sup>, Benjamin Davies<sup>1</sup>, Anna Jones<sup>1</sup>, Dmitri van Popta<sup>1</sup>, Karen Ousey<sup>3</sup>, John Stephenson<sup>3</sup>

<sup>1</sup>Greater Manchester Neurosciences Centre, Manchester Academic Health Science Centre, Salford Royal NHS Foundation Trust

<sup>2</sup>Faculty of Medical and Human Sciences, Manchester Academic Health Science Centre, The University of Manchester

<sup>3</sup>School of Human and Health Sciences, University of Huddersfield

**CORRESPONDING AUTHOR:**

Ross A. Atkinson, PhD

Honorary Research Associate,

Institute of Inflammation and Repair, Manchester Academic Health Sciences Centre, University of Manchester

Email: [ross.atkinson@manchester.ac.uk](mailto:ross.atkinson@manchester.ac.uk)

25

## 26 **SUMMARY**

### 27 **Background**

28 Patients with metastatic spinal tumours have a limited prognosis. Surgical complications which may  
29 result in prolonged hospitalisation or readmission are highly undesirable. Surgical site infection (SSI)  
30 is one such complication which can, in extreme cases, lead to death.

### 31 **Aim**

32 To assess the impact of SSI on patient survival after surgery for spinal metastases.

### 33 **Methods**

34 Demographic, operative and survival data were collected on 152 patients undergoing surgery for  
35 spinal metastasis at a large UK tertiary referral centre. American Society of Anesthesiologists (ASA)  
36 grade and the Revised Tokuhashi Score (RTS) were determined as measures of health status and  
37 prognosis, respectively, at baseline. A semi-parametric Cox proportional hazards survival analysis  
38 was used to assess the relationships between covariates and survival.

### 39 **Findings**

40 Seventeen patients (11.2%) experienced SSI. Overall, median survival time from operation was 262  
41 days (95% CI: 190-334 days) and 12 month survival was 42.1%. RTS ( $p<0.001$ ; hazard ratio 0.82; 95%  
42 confidence interval: 0.76-0.89) and ASA grade ( $p=0.028$ ; hazard ratio 1.37; 95% confidence interval:  
43 1.03-1.82) were significantly associated with survival, with better survival found in patients with  
44 higher RTS and lower ASA scores. Infection status was of substantive importance, with better  
45 survival in those without SSI ( $p=0.075$ ).

### 46 **Conclusion**

47 Twelve month survival in patients undergoing surgery for spinal metastasis is approximately 42%.  
48 RTS and ASA scores can be used as indicators of patient survival either in combination or  
49 individually. While SSI has some negative impact on survival, a larger study sample would be needed  
50 to confirm whether this is statistically significant.

### 51 **Key Words**

52 Metastasis; Spine; Surgical site infection; Survival.

53

## 55 INTRODUCTION

56 Surgical intervention for patients with metastatic spinal tumours is indicated in patients with pain,  
57 instability, or neurological compromise who have a life expectancy exceeding 3 months, according to  
58 the National Institute of Health and Care Excellence (NICE) Clinical Guideline 75<sup>1</sup>. Surgery aims to  
59 prevent or relieve pain, and symptoms associated with mechanical instability and neurological  
60 compromise<sup>2,3</sup>. Complications which jeopardise the success of the operation are therefore highly  
61 undesirable. Surgical site infection (SSI) is one such complication, being the third most commonly  
62 occurring healthcare-associated infection (HAI) in England (accounting for approximately 16%). SSI  
63 has a significant impact on the management of patients undergoing spinal surgery, the length of  
64 time they spend in hospital, and how much money is spent on additional treatment<sup>4</sup>.

65 Although there is little evidence available to demonstrate a direct link between SSI and mortality  
66 specifically in spinal tumour patients, it has been suggested that those suffering SSI as a result of  
67 several types of procedure are overall twice as likely to die and 60% more likely to spend time in the  
68 intensive care unit<sup>5</sup>. Determining the risk of SSI and its potential impact on mortality in patients  
69 undergoing specialist surgical treatment will further highlight the importance of this complication  
70 and the need to implement preventative measures.

71 A significant proportion of patients presenting with spinal metastases are unaware that they have a  
72 primary tumour, with symptoms of spinal cord compression being the first indication of the disease.  
73 Others, however, are known to be suffering from cancer and receiving treatment. Baseline health  
74 status for those undergoing surgery can vary substantially between patients. The Revised Tokuhashi  
75 Score (RTS) is a published system recommended by NICE as a tool to determine eligibility for surgery,  
76 as it is recognised as being able to accurately predict survival<sup>6</sup>. However, some healthcare  
77 providers do not use the RTS to strictly determine which patients should be offered an operation  
78 because not all have a bone scan or a staging computed tomography (CT) pre-operatively, and  
79 because of the perceived palliative benefits of surgery even in those who might have a life  
80 expectancy less than three months. Generally, the RTS suggests that those with a score of 8 or less  
81 have a predicted survival of up to 6 months, meaning that conservative or palliative treatments are  
82 indicated. Those with scores of 12 to 15 have a predicted survival of more than 12 months, meaning  
83 that excisional surgery may be appropriate. Those with a RTS from 9 to 11 are more often suitable  
84 for palliative surgery rather than excisional surgery. Given that RTS appears to be useful in  
85 predicting outcomes for patients with spinal metastasis, this score could be used to standardise  
86 patients at baseline (i.e. the time of their operation) when investigating whether there is an  
87 association between SSI and mortality. Similarly, the American Society of Anesthesiologists (ASA)

88 grade is a recognised scoring system which can be used to determine patient fitness for surgery <sup>7</sup>.  
89 ASA grade is relatively more simple than the RTS to determine, and is routinely recorded prior to  
90 surgery in the majority of cases. Its use is also recommended by NICE in determining the  
91 appropriateness of surgery for patients with spinal tumours.

92 Therefore, the aim of this study was to assess the impact of SSI on patient survival following surgery  
93 for spinal metastatic tumours, after controlling for baseline fitness using both RTS and ASA grades.

94

95

## METHODS

This was a sub-study of an ethically approved case note review of all adult patients (aged  $\geq 18$  years) who had undergone surgical treatment for spinal metastatic tumours at Salford Royal NHS Foundation Trust (SRFT) between 1<sup>st</sup> January 2009 and 31<sup>st</sup> December 2012<sup>8</sup>. Demographic (age, sex), operative (date of operation, type of procedure, presence or absence of SSI) and survival data (date of death, if applicable) were collected. Final follow up assessment was conducted on 15<sup>th</sup> July 2014. In addition, the RTS was determined retrospectively using available medical records and ASA grade was obtained from the surgical documentation. RTS and ASA were determined to give an indication of health status at baseline (i.e. the time of surgery).

### *Definition of SSI*

The presence or absence of a SSI (superficial or deep) was defined using the criteria set out by Public Health England<sup>9</sup>, which is largely based on the definitions published by the Centers for Disease Control and Prevention (CDC) and the work of Horan et al.<sup>10</sup>. SSIs were classified by the SSI surveillance nurse for the neurosurgery department, as per standard routine for the reporting of SSIs through the hospital SSI Surveillance Service.

### *Data Collection*

Data were collected from existing patient case notes and associated medical records (e.g. medical images) and were anonymised prior to analysis; no contact with patients or relatives was required for additional data collection. RTS was calculated based on relevant clinically available data; ideally, RTS would be determined pre-operatively in order that this score can be used to assist in assessing patients' suitability for surgery. However, RTS is not formally recorded routinely at our institution and so this score was determined retrospectively based on the method described by Tokuhashi et al.<sup>6</sup> where possible, with the exception of instances where the number of extra-spinal bony metastases was unavailable from bone scintigraphy or magnetic resonance imaging (MRI). In these cases, staging CT was used. ASA grade was recorded directly from the surgical pathway documentation. All patients were followed up at at least one year post-surgery. Data relating to infection status and survival were collected at this time point.

### *Statistical Analysis*

Overall survival was assessed using the Kaplan-Meier method. A semi-parametric Cox proportional hazards survival (time-to-event) analysis was undertaken to assess the relationships between covariates and survival.

## RESULTS

A total of 152 patients (77 females and 75 males) underwent surgery for spinal metastasis over the four year study period. Mean age at operation was 60.5 years (SD 12.9 years). Seventeen patients (11.2%) experienced SSI (14 superficial and 3 deep). At the time of last follow up, 117 patients had died. Median survival time from operation for the whole cohort was 262 days (95% CI: 190-334 days). This equates to 42.1% at 12 months, and 19.6% at five years. Median survival time for patients experiencing SSI was 135 days (95% CI: 62-208 days), and for those without infection, 276 days (95% CI: 183-369 days).

The assumption of proportional hazards was found to be tenable, and measures of patient fitness were not excessively correlated. The Cox analysis found both RTS and ASA score to be significantly associated with survival (Table 1), with better survival found in patients with higher RTS and lower ASA scores.

Direction of approach was not considered as a candidate factor due to only one patient with an SSI experiencing an anterior approach, and less than 10% of patients in total experiencing an anterior approach. Hence direction of approach does not adequately distinguish between either cases, or between controls and was thus unsuitable as a candidate variable. All cases (infection and non-infection) were instrumented and so this too was not included in the analysis.

[INSERT TABLE 1]

Survival curves for RTS and ASA are shown in Figures 1 and 2. Each additional point on the RTS scale was associated with an 18% lowered hazard of death ( $p<0.001$ ). Each additional point on the ASA scale is associated with a 37% raised hazard of death ( $p=0.028$ ).

[INSERT FIGURES 1 AND 2]

Infection status was found to be of substantive importance, with better survival shown by those without SSI ( $p=0.075$ ) (Figure 3). Age at the time of surgery was not substantively related to survival ( $p=0.299$ ).

[INSERT FIGURE 3]

## DISCUSSION

The results of this study suggest that the median length of survival from the date of surgery for spinal metastases is approximately 8.6 months. This is comparable to figures previously reported in the literature <sup>11, 12</sup>. Twelve month survival in the present cohort was 42.1% overall (29.4% in those with SSI and 43.7% in those without SSI).

As expected, both ASA grade and RTS predicted survival effectively in patients in this study. When using these scores to control for baseline health status, patients experiencing SSI survived on average half as long as those without SSI (though the association between infection and survival was substantive). The low number of cases represented in this study may be a limiting factor and a reason why this association was not observed to be significant. Nevertheless, this study describes the contemporary SSI rate in this patient group at a large UK specialist spinal centre, and is one of the first to demonstrate a relationship between SSI and mortality in this type of patient.

It has been estimated that between 38 and 75% of deaths in patients with SSI are attributable to infection itself <sup>13, 14</sup>, with SSI being an independent predictor of mortality. Other studies have suggested that the type of surgery, and whether SSIs are deep/organ space have some bearing on the level of contribution of infection towards death <sup>15</sup>. While this seems logical, it is impossible in this study to indicate whether the type of infection (superficial or deep) had any effect on survival, given that only three cases were documented as being deep. A larger case series would be needed to determine whether additional factors such as this played an important role in affecting survival outcome.

It is suggested that there are approximately 4,000 cases of metastatic spinal cord compression diagnosed each year in England and Wales <sup>1</sup>, though this is likely to be an underestimate. Surgery is generally undertaken as a palliative measure to relieve pain and stabilise the spine to prevent further neurological damage. Estimating that only around 20% of these will undergo surgery (Richards, personal communication), and based on the 11.2% infection rate demonstrated in this study, 90 patients per year could be at risk of developing SSI. Given the differential in life expectancy demonstrated in this study between patients with and without SSI (141 days shorter in SSI patients), theoretically, this equates to approximately 12,600 days (34.5 years) of life lost which relates to the onset of SSI for just those treated surgically over the period of one year. This, coupled with the distressing consequences of infection (prolonged in-patient hospitalisation, isolation away from home, additional treatments) and the inevitable economic costs associated with SSI <sup>4</sup>, can be exceedingly frustrating for a service under pressure to provide better care at a lower cost. Scrupulous peri-operative practice and the standardisation of effective processes in the operating theatre may go some way to improving outcomes in terms of SSI rates <sup>16-18</sup>. In addition, effective



surgical work-up and the notion of ‘prehabilitation’ in the pre-operative period may better prepare some elective spinal surgery patients physiologically to improve some outcomes<sup>19</sup>. However, there is little evidence to support this approach in reducing post-operative complication rate. Furthermore, in urgent cases – such as those needing surgery for spinal tumours – this window of opportunity is generally not available to the care team, meaning that extra preventative measures are highly desirable and clearly warranted. This is especially so in cancer patients given that they are at increased risk of SSI, due largely to their immunosuppressed state<sup>20, 21</sup>.

The results provided by this study may be useful during the consenting process for spinal surgery. The appropriateness and quality of information patients receive prior to their operation, about the procedure and its associated risks are of utmost importance when ensuring their decisions are fully informed and their expectations managed. Thus, patients have a right to know about the implications of SSI. Despite this, it is evident from previous studies that many patients are poorly informed about SSIs; some are unable to recognise the signs and symptoms of an infection or unaware of the causes or the risk factors of SSI<sup>22</sup>. This low level of awareness about SSI, coupled with the potential for such devastating effects on quality of life and clinical outcome indicates that greater attention should be paid to this complication across the board. Statistics relating to individual departments, such as those presented in this study, could go some way to educating both staff and patients of the institution, in an effort to emphasise risk factors for SSI, what can be done to combat them, and the importance of their consequences. It is hoped this would then translate into the adoption of processes – on the part of both patient and care team – which drive down the incidence of SSI.

## **CONCLUSIONS**

One-year survival in patients undergoing surgery for spinal metastases is approximately 42%. Either or both of RTS or ASA scores can be used as reliable indicators of survival in these patients. While SSI has some negative impact on survival, a larger study sample would be needed to confirm that this is a statistically significant association. The evidence provided by this study may raise awareness of the importance of SSI as a complication of surgery for spinal metastasis.

## **ACKNOWLEDGEMENTS**

This work was supported by a small grant from the Healthcare Infection Society. We would like to thank Lena Richards, Mr Brad Williamson and Dr Chinari Subudhi for their support with this work.

## 228 REFERENCES

- 229 1. NICE, *Metastatic spinal cord compression: diagnosis and management of patients at risk of*  
 230 *or with metastatic spinal cord compression*. 2008, National Collaborating Centre for Cancer:  
 231 Cardiff.
- 232 2. Klimo P, Jr. and Schmidt MH. Surgical management of spinal metastases. *Oncologist*, 2004.  
 233 **9**(2): p. 188-96.
- 234 3. Laufer I, Sciubba DM, Madera M, *et al*. Surgical management of metastatic spinal tumors.  
 235 *Cancer Control*, 2012. **19**(2): p. 122-8.
- 236 4. Urban JA. Cost analysis of surgical site infections. *Surg Infect (Larchmt)*, 2006. **7 Suppl 1**: p.  
 237 S19-22.
- 238 5. Kirkland KB, Briggs JP, Trivette SL, Wilkinson WE and Sexton DJ. The impact of surgical-site  
 239 infections in the 1990s: attributable mortality, excess length of hospitalization, and extra  
 240 costs. *Infect Control Hosp Epidemiol*, 1999. **20**(11): p. 725-30.
- 241 6. Tokuhashi Y, Matsuzaki H, Oda H, Oshima M and Ryu J. A revised scoring system for  
 242 preoperative evaluation of metastatic spine tumor prognosis. *Spine (Phila Pa 1976)*, 2005.  
 243 **30**(19): p. 2186-91.
- 244 7. Dripps RD, Lamont A and Eckenhoﬀ JE. The role of anesthesia in surgical mortality. *JAMA*,  
 245 1961. **178**: p. 261-6.
- 246 8. Atkinson RA, Stephenson J, Jones A and Ousey KJ. An assessment of key risk factors for  
 247 surgical site infection in patients undergoing surgery for spinal metastases. *Journal of Wound*  
 248 *Care*, 2016. **In Press**.
- 249 9. PHE, *Protocol for the surveillance of surgical site infection: surgical site infection surveillance*  
 250 *service*. 2013, Public Health England: London.
- 251 10. Horan TC, Gaynes RP, Martone WJ, Jarvis WR and Emori TG. CDC definitions of nosocomial  
 252 surgical site infections, 1992: a modification of CDC definitions of surgical wound infections.  
 253 *Infect Control Hosp Epidemiol*, 1992. **13**(10): p. 606-8.
- 254 11. Boogerd W and van der Sande JJ. Diagnosis and treatment of spinal cord compression in  
 255 malignant disease. *Cancer Treat Rev*, 1993. **19**(2): p. 129-50.
- 256 12. Moon KY, Chung CK, Jahng TA, Kim HJ and Kim CH. Postoperative survival and ambulatory  
 257 outcome in metastatic spinal tumors : prognostic factor analysis. *J Korean Neurosurg Soc*,  
 258 2011. **50**(3): p. 216-23.
- 259 13. Astagneau P, Rioux C, Golliot F, Brucker G and Group INS. Morbidity and mortality associated  
 260 with surgical site infections: results from the 1997-1999 INCISO surveillance. *J Hosp Infect*,  
 261 2001. **48**(4): p. 267-74.
- 262 14. Awad SS. Adherence to surgical care improvement project measures and post-operative  
 263 surgical site infections. *Surg Infect (Larchmt)*, 2012. **13**(4): p. 234-7.
- 264 15. Coello R, Charlett A, Wilson J, *et al*. Adverse impact of surgical site infections in English  
 265 hospitals. *J Hosp Infect*, 2005. **60**(2): p. 93-103.
- 266 16. Humphreys H. Preventing surgical site infection. Where now? *J Hosp Infect*, 2009. **73**(4): p.  
 267 316-22.
- 268 17. Crolla RM, van der Laan L, Veen EJ, *et al*. Reduction of surgical site infections after  
 269 implementation of a bundle of care. *PLoS One*, 2012. **7**(9): p. e44599.
- 270 18. Johnson B, Starks I, Bancroft G and Roberts PJ. The effect of care bundle development on  
 271 surgical site infection after hemiarthroplasty: an 8-year review. *J Trauma Acute Care Surg*,  
 272 2012. **72**(5): p. 1375-9.
- 273 19. Nielsen PR, Jorgensen LD, Dahl B, Pedersen T and Tonnesen H. Prehabilitation and early  
 274 rehabilitation after spinal surgery: randomized clinical trial. *Clin Rehabil*, 2010. **24**(2): p. 137-  
 275 48.
- 276 20. Olsen MA, Nepple JJ, Riew KD, *et al*. Risk factors for surgical site infection following  
 277 orthopaedic spinal operations. *J Bone Joint Surg Am*, 2008. **90**(1): p. 62-9.

- 278 21. Demura S, Kawahara N, Murakami H, *et al.* Surgical site infection in spinal metastasis: risk  
279 factors and countermeasures. *Spine (Phila Pa 1976)*, 2009. **34**(6): p. 635-9.
- 280 22. Tanner J, Padley W, Davey S, Murphy K and Brown B. Patient narratives of surgical site  
281 infection: implications for practice. *J Hosp Infect*, 2013. **83**(1): p. 41-5.

282