



University of HUDDERSFIELD

University of Huddersfield Repository

Williams, Graham, Uchimoto, Mari, Coult, Natalie, World, Damien and Beasley, Emma

Body fluid mixtures; resolution using forensic microRNA analysis

Original Citation

Williams, Graham, Uchimoto, Mari, Coult, Natalie, World, Damien and Beasley, Emma (2013) Body fluid mixtures; resolution using forensic microRNA analysis. In: The 25th World Congress of the International Society for Forensic Genetics, 2nd-7th September 2013, Melbourne, Australia.

This version is available at <http://eprints.hud.ac.uk/id/eprint/19360/>

The University Repository is a digital collection of the research output of the University, available on Open Access. Copyright and Moral Rights for the items on this site are retained by the individual author and/or other copyright owners. Users may access full items free of charge; copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational or not-for-profit purposes without prior permission or charge, provided:

- The authors, title and full bibliographic details is credited in any copy;
- A hyperlink and/or URL is included for the original metadata page; and
- The content is not changed in any way.

For more information, including our policy and submission procedure, please contact the Repository Team at: E.mailbox@hud.ac.uk.

<http://eprints.hud.ac.uk/>

Introduction

In many cases, especially in sexual assaults, stains can be a combination of different body fluids; for example, semen and vaginal material or blood and saliva. Whilst there are many strategies for the successful resolution of DNA mixtures, there are currently no analytical methods for resolving mixtures of body fluids. RNA analysis is becoming increasingly common place, utilising mRNA and microRNA (miRNA) for body fluid identification purposes. Provisional research has been conducted on using RNA analysis to resolve body fluid mixtures; with Hartevelde *et al* [1] working on mRNA and Uchimoto *et al* [2] working on miRNA. The aim of this study is to further expand on the Uchimoto study [2] and assess whether the major-minor contributors are maintained post-analysis and whether or not the major body fluid can be associated with the major DNA profile and the same for the minor contributor.

Sample acquisition, extraction, and mixture preparation

Blood and saliva samples were acquired from a number of volunteers with their informed consent and ethical approval. Blood samples were acquired using a finger-prick method, and saliva samples were collected via buccal swabs. Both sets of samples were extracted using the buccal protocol of the QIAamp DNA Mini extraction kit (Qiagen, UK) as per the Omelia *et al* study [3]. Blood and saliva extracts were used to prepare mixed sample by volume. Two sets of mixing ratios of 1:1, 5:1, and 10:1 were prepared; one with blood as the major contributor and one with saliva as the major contributor.

sIRT-qPCR

All prepared samples then underwent stem-loop reverse transcription (sIRT) [4] before undergoing quantitative PCR (qPCR). A Veriti thermocycler and a Fast 7500 Real-Time PCR machine were used for analysis along with reverse transcription kits, Taqman master mixes and miRNA assays (Life Technologies, UK). Markers miR-451 and miR-205 were targeted for blood and saliva, respectively. The small nuclear RNA molecule RNU-44 was used as the endogenous control.

Controls

PCR and reverse transcription negative controls were included in this study, along with single source controls. No amplification was detected in the negative controls for this study. A 1:1 mixed body fluid sample was included as a control.

Data analysis

Δ Ct values were derived from CtRNU44-Ctmarker. These values were used for subsequent data analysis. For analysis of mixed samples, a normalising formula was derived from the 1:1 control as per the Uchimoto study. This formula was applied to all samples within the study.

DNA profiling

Aliquots of the prepared mixed samples underwent standard STR analysis, using a Quantiplex kit (Qiagen, UK), NGM SElect Kit (Life Technologies, UK) and the ABI 3130 Genetic Analyser (Life Technologies, UK) with the data analysed by GeneMapper V3.2 Software.

Results

DNA profiling

The aliquots from the prepared mixed samples underwent standard DNA profiling and in all cases mixed DNA results were obtained that matched the corresponding components of the DNA profiles of the donors. In all cases, the overall mixing ratio of the DNA results were comparatively maintained post analysis.

Where the qPCR data indicated that the major contributor was blood, the DNA profile obtained was also from the blood and the same was true of the saliva. This was true of all samples; thus giving a correct body fluid-DNA attribution rate of 100%.





Fig 1; A chart indicating the relative expression of miR-451 and miR-205 in mixed body fluid samples.

The data shown in Figure 1 indicates that the mixing ratios do not appear to be maintained post analysis. For example, the 5:1 and 10:1 mixing ratios appears to indicate that, even though the major and minor contributors are correctly identified, there is no substantial difference between them.

This research was funded internally by the University of Huddersfield. There are no conflicts of interest



The image to the left shows the top two panels of an NGM SElect STR EPG. The sample profiled is a mixture of blood and saliva from two different females. The mixing ratio by volume was 5:1 with blood as the major. In this sample, the major DNA profile identified was from the blood donor. This EPG is a typical example.