



University of HUDDERSFIELD

University of Huddersfield Repository

Camp, Jason, Bousfield, Thomas W., Mistry, Liam and Mapesa, Kopano

Synthesis of ureas in the bio-alternative solvent Cyrene

Original Citation

Camp, Jason, Bousfield, Thomas W., Mistry, Liam and Mapesa, Kopano (2017) Synthesis of ureas in the bio-alternative solvent Cyrene. *Green Chemistry*. pp. 2123-2128. ISSN 1463-9262

This version is available at <http://eprints.hud.ac.uk/id/eprint/31795/>

The University Repository is a digital collection of the research output of the University, available on Open Access. Copyright and Moral Rights for the items on this site are retained by the individual author and/or other copyright owners. Users may access full items free of charge; copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational or not-for-profit purposes without prior permission or charge, provided:

- The authors, title and full bibliographic details is credited in any copy;
- A hyperlink and/or URL is included for the original metadata page; and
- The content is not changed in any way.

For more information, including our policy and submission procedure, please contact the Repository Team at: E.mailbox@hud.ac.uk.

<http://eprints.hud.ac.uk/>

Synthesis of Ureas in the Bio-alternative Solvent Cyrene

Liam Mistry, Kopano Mapesa, Thomas W. Bousfield and Jason E. Camp*

Department of Chemical Sciences, University of Huddersfield
Queensgate, Huddersfield, HD1 3DH, United Kingdom

Abstract – Cyrene as a bio-alternative solvent: a highly efficient, waste minimizing protocol for the synthesis of ureas from isocyanates and secondary amines in the bio-available solvent Cyrene is reported. This method eliminated the use of toxic solvents, such as DMF, and established a simple work-up procedure for removal of the Cyrene, which led to a 28-fold increase in molar efficiency versus industrial standard protocols.

Introduction - Ureas are an important class of compound that have been exploited in a number of fields, such as pharmaceuticals, agrochemicals and materials science. In the pharmaceutical sector the utility of ureas can be observed in the variety of biological activities that they exhibit.¹ For example, ureas are found in anti-malaria compounds,² hepatitis C protease inhibitors,³ HIV-protease inhibitors,⁴ anti-obesity therapeutics,⁵ anticancer agents,⁶ antibiotics,⁷ anticonvulsants⁸ and antimicrobials⁹ (Figure 1). Furthermore, ureas themselves are increasingly exploited as catalysts,¹⁰ ligands,¹¹ reagents,¹² solvents¹³ and substrates¹⁴ for a multitude of synthetic transformations.




Figure 1. Biologically active ureas

Due to their importance a variety of methods have been developed for the synthesis of ureas,¹⁵ with a recent emphasis on developing greener, more sustainable processes.^{16,17} One of the most common ways to synthesize ureas is from the reaction of isocyanates with amines. For example, a series of phenyl isocyanates **1** was reacted with *N*-aryl-*NH*-piperazine **2** to give a series of ureas **3** that was screened for inhibition of platelet-derived growth factor receptor phosphorylation (Figure 2, eq. 1).¹⁸ In this typical example using DMF as solvent, a variety of ureas **3** were formed in good yields, though extended reaction times and extensive work-up / purification procedures needed to be employed. Whilst this reaction can be carried out in a variety of different solvents / reaction media, nearly 80% of the reactions in the literature use DMF or halogenated solvents.¹⁹ There are strong environmental, safety and regulatory pressures to minimize the use of solvents that are designated as being high risk. Replacement for dipolar aprotic solvents,²⁰ such as problematic solvents DMF, NMP and DMAc, was recently highlighted as a priority within the pharmaceutical industry.²¹ In turn, companies and academic researchers have put significant amounts of time, effort and expense into the development of more sustainable chemical processes that do not rely on high risk solvents.^{22,23} A possible alternative to DMF and other polar aprotic solvents is Cyrene (**7**, dihydrolevoglucosenone, Figure 2).²⁴ Cyrene is synthesised in a two-step process from waste cellulose.^{25,26} Importantly, it has similar physical properties to other dipolar aprotic solvents, including DMF. The viability of Cyrene as a DMF replacement was recently demonstrated by Watson *et al.* in palladium mediated cross-coupling reactions.²⁷ In addition, Cyrene^{28,29} was shown to be an effective solvent in S_N2 and S_NAr reactions²⁴ as well as for the synthesis of MOFs.³⁰ Interestingly, the use of Cyrene as a solvent for the addition of a nucleophile to a carbonyl has not been investigated despite the fact that these processes are known to be promoted by polar aprotic solvents.³¹ Herein, we present the development and scope of the use of the bio-available solvent Cyrene (**7**) for the synthesis of ureas **6** from the reaction of isocyanates **4** with amines **5** (Figure 2, eq. 2). An emphasis was placed on minimizing the amount of waste produced during the work-up, isolation and purification of the product.





Figure 2. Reaction of isocyanates and amines in DMF and Cyrene.

Results and Discussion - To begin the investigation, the reaction of phenylisocyanate (**4a**) and pyrrolidine (**5a**) in Cyrene (**7**) was investigated (Scheme 1). Whilst the optimization of the reaction was straightforward, the isolation of the pure urea was more problematic. In order to minimize by-products, the reagents were added together at 0 °C and allowed to warm to rt for 1 h. This protocol allowed for the development of a general procedure for both electron rich and electron deficient isocyanates (*vide infra*). Initially, the product was isolated by the addition of water and dichloromethane, liquid-liquid extraction, washing with water / brine, drying over magnesium sulphate and purified by column chromatography (hexanes/ethyl acetate). This extended procedure resulted in a high yield of the desired urea **6a**, 90%, but also led to a significant amount of waste in the form of organic solvents and contaminated aqueous waste. In addition, the Cyrene was found to co-elute with the product and additional column purification was sometimes required in order to get pure compounds. In order to minimize the use of organic solvents as well as the amount of waste produced in the isolation of the product, it was decided to examine alternative work-up procedures. The use of various acid/base work-up procedures all led to decomposition of both the urea and Cyrene. Finally, it was found that the addition of water to the Cyrene solution at the conclusion of the reaction resulted in precipitation of the desired urea. Filtration and washing with water gave analytically pure *N*-phenylpyrrolidine-1-carboxamide (**6a**) in a synthetically useful yield of 80% (*cf.* Scheme 2).




Scheme 1. Optimization of the synthesis of urea **6a** in Cyrene.

In order to access the efficiency of the two protocols, molar efficiency calculations were undertaken using the method of Watson *et al.*³² in which:


$$\text{molar efficiency (Mol. E\%)} = [\text{moles product} / \text{moles starting material} + \text{additives} + \text{catalysts} + \text{solvents}] \times 100$$

The Mol. E% was calculated for each step of the process and the total molar efficiency ($\text{molE}_{\text{total}}$) is the multiplication of these values. Comparison of the molar efficiency of our original process, which was based on the industrial standard protocol, versus the optimized water washing method showed that use of the latter gave a 28-fold increase in molar efficiency (Figure 3).³³ Similar water addition work-up procedures have been employed to help minimize the E-factors.³⁴ This simple procedure also removed all of the Cyrene from the product, which has been shown to be one of the key challenges of using this solvent.²⁷ Importantly, this method allowed for elimination of all non-bioderived solvents from the process with only Cyrene and water required.

Industrial Standard Aqueous Work-up**work-up / isolation / purification**

$$\text{molE\%} = \frac{\text{mmol total out} = 0.48 \text{ mmol}}{\text{mmol total in} = 2270.9 \text{ mmol}} \times 100 = 2.1 \times 10^{-2}$$


$$\Delta \text{ molE\%} = 28.4$$

Optimized Water Wash Procedure

$$\text{molE\%} = \frac{\text{mmol total out} = 0.40 \text{ mmol}}{\text{mmol total in} = 66.5 \text{ mmol}} \times 100 = .60$$


Figure 3. Comparison of molar efficiency for the synthesis of ureas in Cyrene

With the optimized reaction and isolation procedure in hand the scope of the reaction of amines with phenyl isocyanates (**4a**) in Cyrene was examined (Scheme 2). A variety of secondary cyclic **5a-d** and acyclic amines **5e-n** were subjected to the reaction conditions.³⁵ It was found that in addition to pyrrolidine (**5a**), cyclic amines piperidine (**5b**), morpholine (**5c**) and 1,2,3,4-tetrahydroisoquinoline (**5d**) gave the desired ureas, **6a-d** in good yield. *N,N*-Dialkyl amines **5e-j** with chain length between 2-8 carbons afforded ureas **6e-j** in good yields. In addition, *N,N*-dicyclohexylamine (**5k**) gave urea **6k** in 87% yield. Next, unsymmetrical secondary amines were subjected to the reaction conditions. Both *N*-methyl-*N*-allylamine (**5l**) and *N*-methylaniline (**5m**) gave the desired ureas, **6l** and **6m**, in good yields when subjected to the standard conditions. Finally, bis-*N*-allylamine (**5n**) afforded 1,1-diallyl-3-phenylurea (**6n**) in excellent yield when reacted with phenyl isocyanate (**4a**).



Scheme 2. Synthesis of ureas **6** from phenyl isocyanate **4a** and amines **5** in Cyrene **7**.

Next, the effects of a substituent on the aryl isocyanates were investigated. Thus, a series of substituted aryl isocyanates **4** were reacted with pyrrolidine (**5a**) to give ureas **6o-u** (Scheme 3). Using the optimized conditions for phenyl isocyanate (**4a**) afforded the desired ureas **6o-u** in moderate to good yield. Having an electron-withdrawing group at the 4-position, such as in 4-fluoro-, 4-chloro- and 4-nitrophenyl isocyanates (**4b-d**), led to increased yields relative to the more electron rich 4-methoxyphenyl isocyanate. The use of 2-fluorophenyl isocyanate gave the desired urea **6t** in a lower yield than the 4-fluorophenyl isocyanate isomer. This result suggests that the steric hindrance imposed by having an *ortho* substituent on the aromatic ring leads to a decrease in yield. Finally, 4-fluorophenyl isocyanate was reacted with *N*-methylaniline under the standard conditions to give 1-methyl-1,3-diphenylurea (**6u**) in good yield.



Scheme 3. Effect of substituent on isocyanate reactivity

In situ ^{19}F NMR was used to compare the rate of the formation of ureas from isocyanates and amines in Cyrene versus the industrial standard solvents DMF and CH_2Cl_2 . The room temperature reaction of 4-fluorophenyl isocyanate (**4b**) and pyrrolidine (**5a**) to give urea **6o** in the three solvents of interest was monitored by NMR spectroscopy. All of the reactions were found to be complete in less than 5 minutes when run at 1 M concentration. The relatively fast rates of these reactions is due to the activating 4-fluoro moiety on the phenyl isocyanate. Importantly, this analysis provided further support for the viability of Cyrene to replace the industrial standard solvents DMF and CH_2Cl_2 in this process.

Conclusions

In summary, a green, mild and efficient approach towards the synthesis of ureas from aryl isocyanates and secondary amines in the bio-alternative solvent Cyrene was developed. Both the scope of the amine nucleophile and the effect of substitution on the aryl isocyanate were investigated. This method provides an important alternative to the current industrial use of DMF and halogenated solvents for this process. Importantly, it eliminates the need for the use of both these toxic solvents as well as any non-bioderived organic solvent. Of even more importance is the fact that simply treating the reaction mixture with water solved one of the key problems associated with the use of Cyrene as a solvent, its removal from the product.

Acknowledgements

The Department of Chemical Sciences at the University of Huddersfield (TWB) and Collaborative Venture Funding from the University of Huddersfield (LM) supported this work. We gratefully acknowledge the donation of Cyrene as well as helpful discussions with Tony Duncan, Dr Warwick Raverty and Dr Ken Van Langenberg from Circa Group, Melbourne, Australia.

¹ P. Sikka, J. K. Sahu, A. K. Mishra and S. R. Hashim, *Med. Chem.*, 2015, **5**, 479–483.

² (a) A. Kumar, D. Paliwal, D. Saini, A. Thakur, S. Aggarwall and D. Kaushik, *Eur. J. Med. Chem.*, 2014, **85**, 147–178. (b) J. N. Domínguez, C. León, J. Rodrigues, N. G. de Domínguez, J. Gut and P. J. Rosenthal, *J. Med. Chem.*, 2005, **48**, 3654–3658.

- ³ W. M. Kazmierski, R. Hamatake, M. Duan, L. L. Wright, G. K. Smith, R. L. Jarvest, J.-J. Ji, J. P. Cooper, M. D. Tallant, R. M. Crosby, K. Creech, A. Wang, X. Li, S. Zhang, Y.-K. Zhang, Y. Liu, C. Z. Ding, Y. Zhou, J. J. Planttner, S. J. Baker, W. Bei and L. Liu, *J. Med. Chem.*, 2012, **55**, 3021–3026.
- ⁴ (a) L. V. Romashov and V. P. Ananikov, *Org. Biomol. Chem.*, 2016, DOI: 10.1039/c6ob01731b. (b) A. C. Myers, J. A. Kowalski and M. A. Lipton, *Bioorg. Med. Chem. Lett.*, 2004, **14**, 5219–5222. (c) D. P. Getman, G. A. DeCrescenzo, R. M. Heintz, K. L. Reed, J. J. Talley, M.L. Bryant, M. Clare, K. A. Houseman, J. J. Marr, R. A. Mueller, M. L. Vazquez, H.-S. Shieh, W. C. Stallings and R. A. Stegeman, *J. Med. Chem.*, 1993, **36**, 288–291.
- ⁵ (a) M. D. McBriar, H. Guzik, R. Xu, J. Paruchova, S. Li, A. Palani, J. W. Clader, W. J. Greenlee, B. E. Hawes, T. J. Kowalski, K. O’Neill, B. Spar and B. Weig, *J. Med. Chem.*, 2005, **48**, 2274–2277. (b) C. Fotsch, J. D. Sonnenberg, N. Chen, C. Hale, W. Karbon and M. H. Norman, *J. Med. Chem.*, 2001, **44**, 2334–2356.
- ⁶ (a) H.-Q. Li, P.-C. Lv, T. Yan and H.-L. Zhu, *Anti-Cancer Agent Med. Chem.*, **9**, 417–480. (b) H. Gurulingappa, M. L. Amador, M. Zhao, M. A. Rudek, M. Hidalgo and S. R. Khan, *Bioorg. Med. Chem. Lett.*, 2004, **14**, 2213–2216.
- ⁷ A. A. Bastian, A. Marcozzi and A. Herrmann, *Nature Chem.*, 2012, **11**, 789–793.
- ⁸ D. Kaufmann, M. Bialer, J. A. Shimshoni, M. Devor and B. Yagen, *J. Med. Chem.*, 2009, **52**, 7236–7248.
- ⁹ D. S. Babu, D. Srinivasulu and V. S. Kotakadi, *Chem. Heterocycl. Comp.*, 2015, **51**, 60–66.
- ¹⁰ (a) J. Saadi and H. Wennemers, *Nature Chem.*, 2016, **8**, 276–280. (b) L. Meazza, J. A. Foster, K. Fucke, P. Metrangolo, G. Resnati and J. W. Steed, *Nature Chem.*, 2012, **11**, 42–47.
- ¹¹ (a) Y. Kununobu, H. Ida, M. Nishi and M. Kanai, *Nature Chem.*, 2015, **17**, 712–717. (b) R. Dalpozzo, *Green Chem.*, 2015, **17**, 3671–3686.
- ¹² (a) D. Kumar, S. R. Vemula and G. R. Cook, *Green Chem.*, 2015, **17**, 4300–4306. (b) A. K. Ghosh and M. Brindisi, *J. Med. Chem.*, 2015, **58**, 2896–2940.
- ¹³ M. Liu, L. Liang, X. Li, X. Gao and J. Sun, *Green Chem.*, 2015, **18**, 2851–2863.
- ¹⁴ T. Nishikata, A. R. Abela, S. Huang and B. H. Lipshutz, *J. Am. Chem. Soc.*, 2010, **132**, 4978–4979.
- ¹⁵ (a) G. S. Kumar, R. A. Kumar, P. S. Kumar, N. V. Reddy, K. V. Kumar, M. L. Kantam, S. Prabhakar and K. R. Reddy, *Chem. Comm.*, 2013, **49**, 6686–6688. (b) H. Lebel and O. Leogane, *Org. Lett.*, 2006, **8**, 5717–5720. (c) A. Boeijen, J. van Ameijde and R. M. L. Liskamp, *J. Org. Chem.*, 2001, **66**, 8454–8462. (d) S.-H Lee, H. Matsushita, B. Clapman and K. D. Janda, *Tetrahedron*, 2004, **60**, 3439–3443; (e) A. Yagodkin, K. Lösckke, J. Weisell and A. Azhayev, *Tetrahedron*, 2010, **66**, 2210–2221.
- ¹⁶ (a) A. Sarkarm S. Santra, S. K. Kundu, A. Hajra, G. V. Zyryanov, O. N. Chupakhin, V. N. Charushin and A. Majee, *Green Chem.*, 2016, **18**, 4475–4525. (b) A. D. Mamuye, S. Monticelli, L. Castoldi, W. Holzer and V. Pace, *Green Chem.*, 2015, **17**, 4194–4197. (c) C. Ruß, F. Ilgen, C. Reil, C. Luff, A. H. Begli and B. König, *Green Chem.*, 2011, **13**, 156–161. (d) C. Wu.; H. Cheng, R. Liu, Q. Wang, Y. Hao, Y. Yu and F. Zhao, *Green Chem.*, 2010, **12**, 188–1816. (e) A. Ion, C. Van Doorslaer, V. Parvulescu, P. Jacobs and D. De Vos, *Green Chem.*, 2008, **10**, 111–116. (f) B. M. Bhanage, S.-i. Fujita, Y. Ikushima and M. Arai, *Green Chem.*, 2004, **6**, 78–80.
- ¹⁷ F. Bigi, R. Maggi and G. Sartori, *Green Chem.*, 2000, **2**, 140–148.
- ¹⁸ K. Matsuno, M. Ichimura, T. Nakajima, K. Tahara, S. Fujiwara, H. Kase, J. Ushiki, N. A. Giese, A. Pandey, R. M. Scarborough, N. A. Lokker, J.-C. Yu, J. Irie, E. Tsukuda, S.-i. Ide, S. Oda and Y. Nomoto, *J. Med. Chem.*, 2002, **45**, 3057–3066.
- ¹⁹ SciFinder search of the reaction of isocyanates and amines revealed that of 317934 reactions, 254600 were run in halogenated solvents or DMF, which is 80% of all reported reactions in the database. Search conducted December 5th 2016.
- ²⁰ For a recent example of the use of bio-alternative solvents in synthesis, see: D. Rasina, A. Lombi, S. Santoro, F. Ferlin and L. Vaccaro, *Green Chem.*, 2016, **18**, 6380–6386.
- ²¹ D. J. C. Constable, P. J. Dunn, J. D. Hayler, G. R. Humphrey, J. L. Leazer Jr., R. J. Linderman, R. J.; Lorenz, K.; Manley, B. A. Pearlman, A. Wells, A. Zaks and T. Y. Zhang, *Green Chem.*, 2007, **9**, 411–420.
- ²² For recent publications on the use of more sustainable solvents and solvent selection guides, see: (a) D. Prat, A. Wells, J. Hayler, H. Sneddon, C. R. McElroy, S. Abou-Shehada and P. J. Dunn, *Green Chem.* 2016, **18**, 288–296; (b) Fergal, P. Byrne, S. Jin, G. Paggiola, T. H. M. Petchey, J. H. Clark, T. J. Farmer, A. J. Hunt, C. R. McElroy and J. Sherwood, *Sustain. Chem. Process*, 2016, 4:7. DOI 10.1186/s40508-016-0051-z; (c) C. M. Alder, J. D. Hayler, R. K. Henderson, A. M. Redman, L. Shukla, L. E. Shuster and H. F. Sneddon, *Green Chem.*, 2016, **18**, 3879–3890; (d) P. M. Murray, F. Bellany, L. Benhamou, D.-K. Bučar, A. B. Tabor and T. D. Sheppard, *Org. Biomol. Chem.*, 2016, **14**, 2373–2384; (e) C. R. McElroy, A. Constantinou, L. C. Jones, L. Summerton and J. H. Clark, *Green Chem.*, 2015, **17**, 3111–3121; (f) F. Pena-Pereira, A. Kloskowski and J. Namieśnik, *Green Chem.* 2015, **17**, 3687–3705; (g) D. Prat, J. Hayler and A. Wells, *Green Chem.*, 2014, **16**, 4546–4551.
- ²³ For recent examples of the use of aqueous or alcoholic solutions in place of high risk solvents from our laboratory, see: (a) R. P. Lester, T. Bham, T. W. Bousfield, W. Lewis and J. E. Camp, *J. Org. Chem.*, 2016, **81**, 12472–12477; (b) S. Kyne and J. E. Camp, *ACS Sustainable Chem. Eng.*, 2017, **5**, 41–48; (c) J. E. Camp, J. J. Dunsford, O. S. G. Dacosta, R. K. Blundell, J. Adams, J. Britton, R.J. Smith, T. W. Bousfield and M. W. Fay, *RSC Adv.*, 2016, **6**, 16115–16131. (d) M. Rezayat, R. K. Blundell, J. E. Camp, D. A. Walsh and W. Thielemans, *ACS Sustainable Chem. Eng.*, 2014, **2**, 1241–1250. (e) J. E. Camp, J. J. Dunsford, E. P. Cannons, W. J. Restorick, A. Gadzhieva, M. W. Fay and R. J. Smith, *ACS Sustainable Chem. Eng.*, 2014, **2**, 500–505. (f) R. P. Lester and J. E. Camp, *ACS Sustainable Chem. Eng.*, 2013, **1**, 545–548.

-
- ²⁴ J. Sherwood, M. De bruyn, A. Constantinou, L. Moitym C. R. McElroy, T. J. Farmer, T. Duncan, W. Raverty, A. J. Hunt, J. H. Clark, *Chem. Comm.*, 2014, **50**, 9650–9652.
- ²⁵ For the synthesis of Cyrene from levoglucosenone, see: G. R. Court, C. H. Lawrence, W. D. Raverty and A. J. Duncan, *US Pat.*, 2012/0111714 A1, 2012.
- ²⁶ For the synthesis of levoglucosenone from cellulose, see: (a) K. Koseki, T. Ebata, H. Kawakami, H. Matsushita, K. Itoh and Y. Noai, *US Pat.*, 5,112,994, 1992; (b) F. Cao, T. J. Schwartz, D. J. McClelland, S. H. Krishna, J. A. Dumesic and G. W. Huber, *Energy Environ Sci.*, 2015, **8**, 1808–1815.
- ²⁷ K. L. Wilson, A. R. Kennedy, J. Murray, B. Greatrex, C. Jaieson, A. J. B. Watson, *Beilstein J. Org. Chem.* 2016, **12**, 2005–2011.
- ²⁸ For the use of Cyrene as a chiral pool reagent, see: (a) K. P. Stockton and B. W. Greatrex, *Org. Biomol. Chem.*, 2016, **14**, 7520–7528.
- ²⁹ For the use of the related levoglucosenone as a chiral pool reagent, see: (a) K. P. Stockton, C. J. Merritt, C. J. Sumbly and B. W. Greatrex, *Eur. J. Org. Chem.*, 2015, 6999–7008; (b) G. F. Giri, M. Danielli, R. A. Marinelli and R. A. Spanevello, *Bioorg. Med. Chem. Lett.*, 2016, **26**, 3955–3957; (c) A. L. Flourat, A. A. M. Peru, A. R. S. Teixeira, F. Brunissen and F. Allais, *Green Chem.*, 2015, **17**, 404–412; (d) A. V. Samet, D. N. Lutov, S. I. Firgang, L. A. Lyssenko and V. V. Semenov, *Tetrahedron Lett.*, 2011, **52**, 3026–3028. (e) M. S. Miftakhov, F. A. Valeev and I. N. Gaisina, *Russ. Chem. Rev.*, 1994, **63**, 869–882; (f) A. V. Bridgwater, D. Meier and D. Radlein, *Org. Geochem.*, 1999, **30**, 1479; (g) Q. Lu, W. M. Xiong, W. Z. Li, Q. X. Guo and X. F. Zhu, *Bioresour. Technol.*, 2009, **100**, 4871. (H) S. H. Krishna, D. J. McClelland, Q. A. Rashke, J. A. Dumesic and G. W. Huber, *Green Chem.* 2017, DOI: 10.1039/C6GC03028A.
- ³⁰ J. Zhang, G. White, M. Ryan, A. J. Hunt and M. J. Katz, *ACS Sustainable Chem. Eng.*, 2016, **4**, 7186–7192.
- ³¹ S. Dabi and A. Zilkha, *Eur. Poly. J.*, 1980, **16**, 475–478.
- ³² McGonagle, F. I.; Sneddon, H. F.; Jamieson, C.; Watson, A. J. B. *ACS Sustainable Chem. Eng.* **2014**, **2**, 523–532.
- ³³ For details on the molar efficiency calculations, see the supporting information.
- ³⁴ B. H. Lipshutz, F. Gallou and S. Handa, *ACS Sustainable Chem. Eng.*, 2016, **4**, 5838–5849.
- ³⁵ For a recent example of the use of secondary amines in *N*-cyanation reactions, see: J. N. Ayers, K. B. Ling and L. C. Morrill, *Org. Lett.*, 2016, **18**, 5528–5531.