

#### **University of Huddersfield Repository**

Ousey, Karen, Gillibrand, Warren P. and Lui, Steve

An investigation into the prevention of blistering in post-operative wounds

#### **Original Citation**

Ousey, Karen, Gillibrand, Warren P. and Lui, Steve (2010) An investigation into the prevention of blistering in post-operative wounds. In: Wounds UK 2010, 15th Nov 2010 to 17th Nov 2010, Harrogate, UK. (Unpublished)

This version is available at https://eprints.hud.ac.uk/id/eprint/9068/

The University Repository is a digital collection of the research output of the University, available on Open Access. Copyright and Moral Rights for the items on this site are retained by the individual author and/or other copyright owners. Users may access full items free of charge; copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational or not-for-profit purposes without prior permission or charge, provided:

- The authors, title and full bibliographic details is credited in any copy;
- A hyperlink and/or URL is included for the original metadata page; and
- The content is not changed in any way.

For more information, including our policy and submission procedure, please contact the Repository Team at: E.mailbox@hud.ac.uk.

http://eprints.hud.ac.uk/

# An investigation into the prevention of blistering in post-operative wounds

AUTHORS: Dr Karen Ousey, Dr Warren Gillibrand, Steve Lui – School of Huddersfield, Queensgate, Huddersfield, West Yorkshire, HD1 3DH, United Kingdom

## INTRODUCTION

Surgical patients are at risk of developing post-operative wound complications including blistering and infection.

Consequently stays in hospital could be lengthened; costs increase and morbidity/ mortality rates can be adversely affected (Tustanowski 2009).

Blistering can cause increased pain, delayed healing and increased susceptibility to wound infection, as the integrity of the skin has been breached (Gupta et al, 2002). There is currently conjecture in the literature as to whether dressing choice has an effect on wound complication rate (Tustanowski 2009). Blistering after hip surgery caused by tape related injuries has been reported as 21.4% (Polatsch et al, 2004). In a quasi-experimental study (Jester et al, 2000) the incidence of blistering, using a variety of dressings, was reported as 13%. In a prospective study of patients undergoing hip or knee surgery, the post-operative blistering rate ranged from 6% to 24% depending on the dressing used (Cosker et al, 2005). The incidence of superficial wound problems, such as skin blistering, is a commonly reported problem, especially in orthopaedic surgery (Wright, 1994; Cosker et al, 2005).

#### METHOD

A rapid appraisal of the literature has been performed to determine current evidence regarding the prevention and management of post operative wound blistering. A prospective cohort study of patients undergoing orthopedic replacement joint surgery is currently being carried out, to assess incidence of wound blistering.

#### RESULTS

A standard systematic search of the literature returned 137 articles related to wounds and healing. Through blind, two person peer review of the abstracts, 9 were identified to have direct relevance to wound blisters and prevention and/ or treatment.

### DISCUSSION

A limited amount of studies have examined the effect of different dressings on wound healing with no conclusive recommendations (Tustanowski 2009). Wound blistering could be associated with a number of factors; movement of the wound site; choice of dressing; tape use; age; gender; type of incision; medications; co-morbidity; costeffectiveness of dressings (Tustanowski 2009). However an overall conclusion has not been reached, other than adherence to the accepted principles of good post-operative wound management, and standard properties of wound dressings. It is argued that calls for further comparative studies of wound dressings will only continue to provide equivocal results. Therefore it may be that the way forward is to achieve consensus between experts and practitioners as to the most clinical and cost effective dressings and postoperative wound management to prevent blistering and other complications.

# CONCLUSION

There is limited data and consensus opinion to the prevention of post operative wound blistering which suggests a need to develop evidence based/best practice guidance to assist practitioners to choose the most appropriate wound dressing that will help to reduce the incidence of post operative blistering.

Morbidity/mortality rates can be adversely affected

Costs increase

Stays in hospital could be lengthened

Surgical patients remain at risk of developing post-operative wound complications including blistering and infection

Blistering can cause increased susceptibility to wound infection, as the integrity of the skin has been breached

Blistering can cause increased pain

Blistering can cause delayed healing

**References**: Cosker T, Elsayed S, Gupta S, Mendonca AD, Tayton KJJ (2005) Choice of dressing has a major impact on blistering and healing outcomes in orthopaedic patients. Journal of Wound Care 14(1): 27-9. Gupta SK, Lee S, Moseley L G (2002) Postoperative wound blistering: is there a link with dressing usage? Journal of Wound Care 11(7): 271-73. Jester R, Russell L, et al (2000) A one hospital study of the effect of wound dressings and other related factors on skin blistering following total hip and knee arthroplasty. Journal of Orthopaedic Nursing 4(2): 71–7. Polatsch DB, Baskies MA, et al (2004) Tape blisters that develop after hip fracture surgery: a retrospective series and a review of the literature. American Journal of Orthopaedics 33(9): 452–6. Tustanowski J (2009) Effect of dressing choice on outcomes after hip and knee arthroplasty: a literature review. Journal of Wound Care, 18, 11, 449-458. Wright M (1994) Hip blisters. Nursing Times 90(16): 86-8